arXiv:cond-mat/0004005v1 [cond-mat.str-el] 1 Apr 2000

Rigorous Schrieffer-Wolff transformation for the Anderson impurity model

Gilles Poirot

Abstract

With the help of computer algebra, I devise an exact unitary transformation for the Anderson impurity model which allows to kill the hybridization term in the slightly simplified case of zero chemical potential. Then I compute explicitly the outcome of this transformation. This is a rigorous version of the well known Schrieffer-Wolff transformation. It should be possible to treat the general case at the price of increased computation time.

1 Introduction

The Anderson impurity model describes a single magnetic impurity coupled to a conduction band of electrons (for a recent review see [1]). The reduced Hamiltonian with chemical potential μ is *formally* given by

$$\begin{aligned}
H' &= H - \mu N && (1)\\
&= H'_0 + (\varepsilon_d - \mu) n_d + U n_{d,\uparrow} n_{d,\downarrow} + g \sum_{\sigma \in \{\uparrow,\downarrow\}} [c^*_\sigma(V) d_\sigma + \text{h.c.}] && (2)\\
H'_0 &= \sum_{\sigma \in \{\uparrow,\downarrow\}} \int dk \, [\varepsilon(k) - \mu] c^*_\sigma(k) c_\sigma(k) && (3)\\
c^*_\sigma(V) &= \int dk \, V(k) c^*_\sigma(k) \; = \; [c_\sigma(V)]^* && (4)\\
n_{d,\sigma} &= d^*_\sigma d_\sigma && (5)\\
n_d &= n_{d,\uparrow} + n_{d,\downarrow} && (6)\\
\|V\|^2_{L^2} &= 1 && (7)
\end{aligned}$$

where in fact one should pass to *particle-hole* representation and define a new H' (by adding a diverging constant) bounded from below in the infinite volume limit. The c's and the d's are standard Fermionic annihilation operators, we assume furthermore

$$\{c_\sigma(k), d_{\sigma'}\} = \{c_\sigma(k), d^*_{\sigma'}\} = 0 \quad (8)$$

(*In the following, I will note H instead of H'.*)

Using a Bethe-ansatz, the model was exactly solved in dimension $d = 1$ with linear dispersion relation (without UV-cutoff) and a constant V [5]. But in the general case, apart from numerical renormalization [3], the common way to study this model consists in trying to eliminate the hybridization term. This was first approximatively done by Schrieffer and Wolff [4]. This Schrieffer-Wolff transformation leads to a renormalization of the energy impurity and of the repulsive interaction. Furthermore it generates an anti-ferromagnetic spin-spin interaction between the impurity and conduction band electrons. Thus, it maps the Anderson impurity model to an effective s-d type model exhibiting Kondo effect.

But this transformation is equivalent to a second order perturbation treatment and its validity is unclear. Furthermore, it becomes singular when the energy of the impurity lies in the conduction band. In particular, it is not at all suitable in the intermediate valence regime (of interest for certain rare earth compounds) where the impurity energy is close to the Fermi level.

The Schrieffer-Wolff transformation was refined by Kehrein and Mielke by the use of infinitesimal unitary transformations [2]. This leads to a smoother result but it still relies on uncontrolled approximations.

In this paper, I want to study the possibility of an *exact* elimination of the hybridization term through unitary transformation with the help of *computer algebra*. As a first step, I will consider the slightly simplified case $\mu = 0$, *i.e.* there are only particles and no holes. In this case, I prove the existence of such a generalized Schrieffer-Wolff transformation and I compute the resulting Hamiltonian as a s-d type model (up to irrelevant terms) with an effective spin-spin interaction which is anti-ferromagnetic in one channel.

Then I explain how the computation would be changed in the general case and how one can then partly use the $\mu = 0$ computation. This suggests that it should be possible to treat also the general case.

2 Main result

Theorem 1 *Let H be as in (2) with $U > 0$, $\mu = 0$ and $\frac{V}{\varepsilon^{1/2}} \notin L^2(\mathbb{R}^d)$, then there exist*

- $(t_1, t_2) \in \mathbb{R}^2$;
- $(f_1, f_2) \in L^2(\mathbb{R}^d)^2$ *with* $\|f_1\|_{L^2}^2 = \|f_2\|_{L^2}^2 = 1$;
- *and some unitary operators*

$$U_1 = \exp\left\{ t_1 \sum_\sigma [c_\sigma^*(f_1) d_\sigma - \text{h.c.}] \right\} \tag{9}$$

$$U_2 = \exp\left\{ t_2 \sum_\sigma [c_\sigma^*(f_2) d_{-\sigma}^* d_{-\sigma} d_\sigma - \text{h.c.}] \right\} \tag{10}$$

such that

$$H_2 = U_2 U_1 H U_1^* U_2^* \tag{11}$$

contains no term linear in c_σ or c_σ^*.

The proof is through brute force. Using the above expressions as ansatz for U_1 and U_2, I compute (with the help of computer algebra) the effect of the two unitary transformations as a function of $(t_1, f_1; t_2, f_2)$ then I impose that the terms linear in c_σ or c^*_σ vanish.

After simplification, I obtain the following expected result

Theorem 2 *H is unitarily equivalent to an s-d type Hamiltonian*

$$\begin{aligned} H_2 &= H_0 + \varepsilon_R n_d + U_R n_{d,\uparrow} n_{d,\downarrow} + K_1 \left[c^*_\uparrow(f_1) c^*_\downarrow(f_1) d_\downarrow d_\uparrow + \text{h.c.}\right] \\ &\quad + \sum_{\substack{\sigma\in\{\uparrow,\downarrow\} \\ i\in\{+,-\}}} \lambda_i c^*_\sigma(\varphi_i) c_\sigma(\varphi_i) - \sum_{\substack{\sigma\in\{\uparrow,\downarrow\} \\ x\in\{AF,F\}}} \lambda_x c^*_\sigma(\varphi_x) c_\sigma(\varphi_x) n_d \\ &\quad +4 \sum_{x\in\{AF,F\}} \lambda_x \vec{S}_c(\varphi_x\otimes\varphi_x) . \vec{S}_d + H_{Irr} \qquad (12) \\ S^{(i)}_c(\varphi\otimes\psi) &= \sum_{(\tau,\tau')\in\{\uparrow,\downarrow\}^2} c^*_\tau(\varphi) \left(\frac{\sigma_i}{2}\right)_{\tau\tau'} c_{\tau'}(\psi), \quad i\in\{x,y,z\} \qquad (13) \\ S^{(i)}_d &= \sum_{(\tau,\tau')\in\{\uparrow,\downarrow\}^2} d^*_\tau \left(\frac{\sigma_i}{2}\right)_{\tau\tau'} d_{\tau'} \qquad (14) \end{aligned}$$

where H_{Irr} is an irrelevant *part (in the Renormalization Group sense) which consists in a (large) number of terms of order at least 3 in c or c^*, and the σ_i are Pauli matrices (thus the $\vec{S}$'s are spin operators).*

Furthermore, for values of the parameters (ε_d, U, g) in the so-called local moment *or* intermediate valence *regimes, one has*

$$\lambda_{AF} > 0 \quad \textit{and} \quad \lambda_F < 0 \qquad (15)$$

i.e. *the resulting spin-spin interaction is anti-ferromagnetic in one channel.*

Remarks

- Let us note that U_1 and U_2 preserve the original *electron number* and *total spin* conservation as well as the $U(1)\times SU(2)$ symmetry. Therefore, the only possible terms linear in c_σ or c^*_σ are

  $$\sum_\sigma [c^*_\sigma(.) d_\sigma + \text{h.c.}] \quad \text{and} \quad \sum_\sigma [c^*_\sigma(.) d^*_{-\sigma} d_{-\sigma} d_\sigma + \text{h.c.}]$$

  this means that I perform a kind of minimal transformation (2 adjustable functions to meet 2 constraints).

- The Schrieffer Wolff transformation amounts to the particular choice

$$\begin{aligned} f_1 &\propto \frac{V}{\varepsilon-\varepsilon_d} \qquad (16) \\ f_2 &\propto \frac{V}{(\varepsilon-\varepsilon_d)\left[\varepsilon-(U+\varepsilon_d)\right]} \qquad (17) \end{aligned}$$

We will see that this is not so bad provided one replaces ε_d and U by their *renormalized value.* Indeed I find

$$\begin{aligned} f_1 &\propto \frac{V}{\varepsilon - \varepsilon_R} && (18)\\ f_2 &\propto a_2 \frac{V}{\varepsilon - \varepsilon_R} + b_2 \frac{V}{(\varepsilon - \varepsilon_R)(\varepsilon + \mu_2)} \quad \text{where } \mu_2 \sim -(U_R + \varepsilon_R) && (19)\end{aligned}$$

- Up to irrelevant terms, one can see that in H_2 the channels $n_d = 1$ and $n_d \in \{0,2\}$ are decoupled. Yet $U_R + \varepsilon_R \sim -\mu_2$ is negative but small (in the local moment regime, I find that $\mu_2 \sim e^{-C|\varepsilon_d|/g^2}$). Thus it is a priori not clear in which channel will be the ground state of H_2.

- Finally an important remark is that strictly speaking eliminating the hybridization term is not enough because there remains a dangerous term in H_2, namely

$$\mathcal{W} = K_1 \left[c^*_\uparrow(f_1)c^*_\downarrow(f_1)d_\downarrow d_\uparrow + \text{h.c.}\right] \tag{20}$$

  The reason is that when one performs some Renormalization Group analysis on H_2, $\mathcal{W}$ combined with some cubic term in H_{Irr} can generate back some terms which are linear in c_σ or c^*_σ. Thus one should perform a third transformation

$$U_3 = \exp\left\{t_3[c^*_\uparrow(f_3)c^*_\downarrow(f_3)d_\downarrow d_\uparrow - \text{h.c.}]\right\} \tag{21}$$

  to kill also this term. This should a priori be possible, one then would have to meet three constraints with three adjustable functions.It is easy to see that there are no other *dangerous* term which could generate some part linear in c_σ or c^*_σ.

  This problem was totally overlooked up to know because people usually do not worry about irrelevant terms. This is not too dramatic in the *local moment* regime since in that case, one has

$$K_1 \sim e^{-C|\varepsilon_d|/g^2} \tag{22}$$

  which means that the terms linear in c_σ or c^*_σ that are generated will in fact remain small at least up to the Kondo Temperature scale.

3 Proof of theorem 1

3.1 Principle of the computation

The unitary transformations that I consider in this paper are of the form $U(t) = e^{t\Gamma}$, and the general problem is, given an observable O, to compute

$$O(t) = e^{t\Gamma} O e^{-t\Gamma} \tag{23}$$

$O(t)$ is solution of the following differential equation

$$\frac{d}{dt}O(t) = e^{t\Gamma}[\Gamma,O]e^{-t\Gamma} = [\Gamma,O](t) \tag{24}$$
$$O(t=0) = O \tag{25}$$

Thus the problem amounts to find a finite (possibly large) set $\{O_1 = O, O_2, \dots, O_n\}$ whose span is stable under commutation with Γ.

$$[\Gamma, O_i] = \sum_j M_{ij} O_j \tag{26}$$

Then one has just to integrate the first order linear equation

$$\frac{dX_t}{dt} = MX_t \quad \text{with} \quad X_{t=0} = (O_1, \dots, O_n) \tag{27}$$

The difficulty comes from the large value of n (for the second transformation, $n \approx 150$) which makes the problem in practical impossible to solve by hand, thus the need for computer assistance.

The computation was done on a Unix workstation running *Mathematica*, using a personal implementation of Fermionic operator algebras. The computation time is hard to assess because computation went in parallel with code writing and furthermore the whole process is not yet automated and requires an heavy human participation.

I can nevertheless estimate a kind of effective computation time (human+computer) which is of order of a few months. This proves that such computations are tractable.

3.2 First Unitary transformation

Let $H_1 = U_1 H U_1^*$, $\alpha_1 = \cos(t_1)$ and $\beta_1 = \sin(t_1)$, this transformation is quite easy to compute since one can check that

$$c_\sigma(f_1) \overset{U_1}{\longmapsto} \alpha_1 c_\sigma(f_1) - \beta_1 d_\sigma \tag{28}$$
$$d_\sigma \overset{U_1}{\longmapsto} \beta_1 c_\sigma(f_1) + \alpha_1 d_\sigma \tag{29}$$
$$c_\sigma(k) \overset{U_1}{\longmapsto} c_\sigma(k) + f_1(k)[(\alpha_1 - 1)c_\sigma(f_1) - \beta_1 d_\sigma] \tag{30}$$

An important simplification coming from the case $\mu = 0$ is the fact that the second transformation does not affect the "$c_\sigma^*(.)d_\sigma$" part. Therefore I can set it to zero at this stage. The constraint reads

$$(\varepsilon + \lambda_1) f_1 = g\frac{\alpha_1}{\beta_1} V \tag{31}$$
$$\lambda_1 = g\frac{\alpha_1}{\beta_1}(1-\alpha_1)\langle f_1, V\rangle - (1-\alpha_1)\langle f_1, \varepsilon f_1\rangle + g\beta_1 \langle V, f_1\rangle - \alpha_1 \varepsilon_d \tag{32}$$

This leads to the following equation for λ_1

$$\lambda_1 = g^2 \left\langle V, \frac{1}{\varepsilon + \lambda_1} V \right\rangle - \varepsilon_d \tag{33}$$

from which one can see that λ_1 is real, just like $\langle V, f_1\rangle$.

Let us consider

$$\varepsilon_d(\lambda) = g^2 \left\langle V, \frac{1}{\varepsilon+\lambda} V \right\rangle - \lambda, \quad \lambda \in (0, +\infty) \tag{34}$$

it is easy to see that this is a strictly decreasing function ranging from $+\infty$ (because $\frac{V}{\varepsilon^{1/2}} \notin L^2$) to $-\infty$. Therefore it admits an inverse function $\lambda_1(\varepsilon_d) > 0$ giving the unique solution to (33).

Since $\lambda_1 > 0$, $f_1 \in L^2(\mathbb{R}^d)$ and the normalization condition fixes α_1^2 and β_1^2.

$$\begin{aligned} \alpha_1^2 &= \frac{1}{1+g^2 M_1^2} \leqslant 1 && (35)\\ M_1^2 &= \left\langle V, (\varepsilon+\lambda_1)^{-2} V \right\rangle && (36) \end{aligned}$$

Finally, choosing α_1 and β_1 to be positive, one obtains.

$$\begin{aligned} H_1 &= H_0 + C_1 d_\uparrow^* d_\downarrow^* d_\downarrow d_\uparrow + C_3[c_\uparrow^*(f_1) c_\downarrow^*(f_1) d_\downarrow d_\uparrow + \text{h.c.}] \\ &\quad + C_5 c_\uparrow^*(f_1) c_\downarrow^*(f_1) c_\downarrow(f_1) c_\uparrow(f_1) + \sum_{\sigma\in\{\uparrow,\downarrow\}} W_\sigma^{(1)} && (37)\\ W_\sigma^{(1)} &= C_2[c_\sigma^*(f_1) d_{-\sigma}^* d_{-\sigma} d_\sigma + \text{h.c.}] - \frac{C_3}{2}[c_\sigma^*(f_1) c_{-\sigma}(f_1) d_{-\sigma}^* d_\sigma + \text{h.c.}] \\ &\quad + C_3 c_\sigma^*(f_1) c_\sigma(f_1) d_{-\sigma}^* d_{-\sigma} + C_4[c_\sigma^*(f_1) c_{-\sigma}^*(f_1) c_{-\sigma}(f_1) d_\sigma + \text{h.c.}] \\ &\quad + C_6[c_\sigma^*(f_1) c_\sigma(\varepsilon f_1) + \text{h.c.}] + C_7 d_\sigma^* d_\sigma + C_8 c_\sigma^*(f_1) c_\sigma(f_1) && (38) \end{aligned}$$

where the C's are given by

$$\begin{aligned} x_1 &= g M_1 && (39)\\ C_1 &= \frac{U}{(1+x_1^2)^2} && (40)\\ C_2 &= x_1 C_1 && (41)\\ C_3 &= x_1^2 C_1 && (42)\\ C_4 &= x_1^3 C_1 && (43)\\ C_5 &= x_1^4 C_1 && (44)\\ C_6 &= -1 + \sqrt{1+x_1^2} && (45)\\ C_7 &= -\lambda_1 && (46)\\ C_8 &= \frac{1}{x_1^2}\left\{(2+x_1^2)\varepsilon_d + (2-x_1^2)\lambda_1 - 2\sqrt{1+x_1^2}\left[\varepsilon_d + \lambda_1(1-x_1^2)\right]\right\} && (47) \end{aligned}$$

3.3 Second Unitary Transform

By analogy with the first transformation, I will look for a function f_2 satisfying

$$\forall\, k \in \mathbb{R}^d, \quad \overline{f_1(k)}\, f_2(k) \in \mathbb{R} \tag{48}$$

The interested reader will find in appendix the differential equations governing this transformation, from this one can reconstruct the integrated flow.

I note

$$\begin{aligned}
\alpha_2 &= \cos(\sqrt{2}\, t_2) && (49)\\
\beta_2 &= \sin(\sqrt{2}\, t_2) && (50)\\
\Gamma_i &= \langle f_i, \varepsilon f_i \rangle \quad i = 1, 2 && (51)\\
\omega_{12} &= \langle f_1, f_2 \rangle \quad (\in \mathbb{R}) && (52)\\
\Gamma_{12} &= \langle f_1, \varepsilon f_2 \rangle \quad (\in \mathbb{R}) && (53)
\end{aligned}$$

Having the $c^*_\sigma(.)d^*_{-\sigma}d_{-\sigma}d_\sigma$ term vanishing leads to the following constraint

$$\begin{aligned}
(\varepsilon + \mu_2)f_2 &= -\omega_{12}C_6(\varepsilon + \mu_1)f_1 && (54)\\
-\omega_{12}C_6\mu_1 &= \sqrt{2}\,\frac{\alpha_2}{\beta_2}C_2 - 2\omega_{12}C_3 - \omega_{12}C_8 - \Gamma_{12}C_6 && (55)\\
\mu_2 &= -\alpha_2(C_1 + C_7) - \Gamma_2(1-\alpha_2) + \sqrt{2}\,\frac{\alpha_2}{\beta_2}\omega_{12}C_2(1-\alpha_2) + \sqrt{2}\,\beta_2\omega_{12}C_2 \\
&\quad -\omega_{12}^2(2C_3 + C_8)(1-\alpha_2) - 2\Gamma_{12}\omega_{12}C_6(1-\alpha_2) && (56)
\end{aligned}$$

Using expression (54) one gets

$$\begin{aligned}
\omega_{12} &= -\omega_{12}C_6\left[1 - (\mu_2 - \mu_1)\left\langle f_1, \frac{1}{\varepsilon + \mu_2} f_1 \right\rangle\right] && (57)\\
\Gamma_{12} &= -\omega_{12}[C_6(\Gamma_1 + \mu_1) + \mu_2] && (58)\\
\Gamma_2 &= \omega_{12}^2 C_6[C_6(\Gamma_1 + \mu_1) + \mu_2 - \mu_1] - \mu_2 && (59)\\
& && (60)
\end{aligned}$$

And from (31), (33) and (35)

$$\Gamma_1 = \frac{\lambda_1 + \varepsilon_d}{g^2 M_1^2} - \lambda_1 \qquad (61)$$

Substituting (58) into (55) yields

$$\begin{aligned}
\alpha_2^2 &= \frac{Z^2}{Z^2 + 2C_2^2} \;<\; 1 && (62)\\
Z &= \omega_{12}\left[-2C_3 - C_8 + C_6\mu_2 + C_6(C_6+1)\mu_1 + \Gamma_1 C_6^2\right] && (63)
\end{aligned}$$

Then I choose ($\theta = \pm 1$)

$$\begin{aligned}
\alpha_2 &= \frac{\theta Z}{\sqrt{Z^2 + 2C_2^2}} && (64)\\
\beta_2 &= \frac{-\theta\sqrt{2}C_2}{\sqrt{Z^2 + 2C_2^2}} && (65)
\end{aligned}$$

With this particular choice, ω_{12} disappears from equation (56) so that I can express μ_1 as a function of μ_2.

$$\mu_1 = -\frac{2C_2^2 + (C_1 + C_7 + \mu_2)\left[-2C_3 - C_8 + C_6(\mu_2 + C_6\Gamma_1)\right]}{C_6(C_6+1)(C_1 + C_7 + \mu_2)} \tag{66}$$

And from the normalization condition, one gets ω_{12} as a function of μ_1 and μ_2.

$$1 = \langle f_2, f_2 \rangle = \omega_{12}^2 C_6^2 \left[-\frac{C_6+2}{C_6} + (\mu_2 - \mu_1)^2 \left\langle f_1, \frac{1}{(\varepsilon + \mu_2)^2} f_1 \right\rangle \right] \tag{67}$$

Finally, substituting (66) into (57), we are left with the following equation

$$\begin{aligned}(1+C_6)^2(C_1 + C_7 + \mu_2) \;&=\; \left\langle f_1, \frac{1}{\varepsilon + \mu_2} f_1 \right\rangle \Big\{ 2C_2^2 \\ &\quad + (C_1 + C_7 + \mu_2)\big[-2C_3 - C_8 + C_6(C_6+2)\mu_2 + C_6^2\Gamma_1 \big] \Big\} \qquad (68)\end{aligned}$$

which after simplification becomes

$$\varphi(\mu_2) = 0 \tag{69}$$

$$\begin{aligned}\varphi(\mu) \;&=\; U + (1+x_1^2)^2\,(\mu - \lambda_1) \\ &\quad + \frac{x_1^2}{1+x_1^2}(\mu - \lambda_1)\big[U - (1+x_1^2)^2\,(\mu - \lambda_1)\big] \left\langle f_1, \frac{1}{\varepsilon + \mu} f_1 \right\rangle \qquad (70)\end{aligned}$$

Lemma 1

The equation $\varphi(\mu) = 0$ has a unique solution μ_2, furthermore $\mu_2 \in (0, \lambda_1)$.

Proof

- First, let us note that φ is a smooth function with

  $$\lim_{\mu \to 0} \varphi(\mu) = -\infty \quad (\text{since } \frac{V}{\varepsilon^{1/2}} \notin L^2) \tag{71}$$

  $$\varphi(\lambda_1) = U > 0 \tag{72}$$

  This implies that there is at least one solution $\mu_2 \in (0, \lambda_1)$.

- For $\mu > \lambda_1$ we have $\varphi(\mu) \geqslant U > 0$ at least as long as $U - (1+x_1^2)^2\,(\mu - \lambda_1) \geqslant 0$, *i.e.* as long as $\mu \leqslant \lambda_1 + U(1+x_1^2)^{-2}$.

- For $\mu > \lambda_1 + U(1+x_1^2)^{-2}$, we note that

  $$\begin{aligned}\varphi(\mu) \;&=\; U + (1+x_1^2)^2\,(\mu - \lambda_1) \\ &\quad - \frac{x_1^2}{1+x_1^2}(\mu - \lambda_1)\big[(1+x_1^2)^2\,(\mu - \lambda_1) - U\big] \left\langle f_1, \frac{1}{\varepsilon + \mu} f_1 \right\rangle \qquad (73) \\ &\geqslant\; U + (1+x_1^2)^2\,(\mu - \lambda_1) \\ &\quad - \frac{x_1^2}{1+x_1^2}(\mu - \lambda_1)\big[(1+x_1^2)^2\,(\mu - \lambda_1) - U\big] \frac{1}{\mu} \qquad (74) \\ &\geqslant\; U + (1+x_1^2)^2\,(\mu - \lambda_1) - x_1^2(1+x_1^2)(\mu - \lambda_1) \qquad (75) \\ &\geqslant\; U + (1+x_1^2)\,(\mu - \lambda_1) \;>\; 0 \qquad (76)\end{aligned}$$

  Thus there are no solution in $[\lambda_1, +\infty)$.

- Finally, for $0 < \mu < \lambda_1$, we have

$$\begin{align}
\varphi'(\mu) &= (1+x_1)^2 + \frac{x_1^2}{1+x_1^2}\big[U + 2(1+x_1^2)^2(\lambda_1-\mu)\big]\left\langle f_1, \frac{1}{\varepsilon+\mu} f_1\right\rangle \nonumber\\
&\quad + \frac{x_1^2}{1+x_1^2}(\lambda_1-\mu)\big[U + (1+x_1^2)^2(\lambda_1-\mu)\big]\left\langle f_1, \frac{1}{(\varepsilon+\mu)^2} f_1\right\rangle \tag{77}\\
&> 0 \tag{78}
\end{align}$$

  therefore there is a unique $\mu_2 \in (0, \lambda_1)$ such that $\varphi(\mu_2) = 0$.

■

Since $\mu_2 > 0$, $f_2 \in L^2(\mathbb{R}^d)$ and I can successively solve (66), (67) and (62).

$$\begin{align}
\mu_1 &= \frac{x_1^2}{(1+x_1^2) - \sqrt{1+x_1^2}}\lambda_1 - \frac{\mu_2}{\sqrt{1+x_1^2}} \nonumber\\
&\quad - \frac{2Ux_1^2}{\big[U - (1+x_1^2)^2(\lambda_1-\mu_2)\big]\left[(1+x_1^2) - \sqrt{1+x_1^2}\right]}(\lambda_1-\mu_2) \tag{79}\\
\omega_{12}^2 &= \left[(\mu_2-\mu_1)^2(\sqrt{1+x_1^2}-1)^2\left\langle f_1, (\varepsilon+\mu_2)^{-2} f_1\right\rangle - x_1^2\right]^{-1} \tag{80}\\
\alpha_2^2 &= \frac{N_{\alpha_2}^2}{D_{\alpha_2}^2} \tag{81}\\
N_{\alpha_2} &= (\mu_2-\mu_1)(\sqrt{1+x_1^2}-1) - x_1^2\left[\frac{2U}{(1+x_1^2)^2} + \lambda_1 - \mu_1\right] \tag{82}\\
D_{\alpha_2}^2 &= N_{\alpha_2}^2 + \frac{2U^2x_1^2}{(1+x_1^2)^4}\omega_{12}^{-2} \tag{83}
\end{align}$$

Furthermore, f_2 is given by

$$f_2 = -\omega_{12}(\sqrt{1+x_1^2}-1)\left(\frac{\varepsilon+\mu_1}{\varepsilon+\mu_2}\right) f_1 \tag{84}$$

This concludes the proof of theorem 1.

4 s-d type Hamiltonian

After simplification, I find that the initial Anderson impurity Hamiltonian H is unitarily equivalent to H_2 whose interacting part, up to irrelevant terms, consists in a potential scattering term and a s-d exchange term.

$$\begin{aligned}
H_2 &= H_0 + \varepsilon_R n_d + U_R n_{d,\uparrow} n_{d,\downarrow} + K_1 \left[c^*_\uparrow(f_1) c^*_\downarrow(f_1) d_\downarrow d_\uparrow + \text{h.c.}\right] \\
&\quad + \sum_\sigma W_\sigma - 4W_{\text{sd}} + H_{Irr}
\end{aligned} \tag{85}$$

$$\begin{aligned}
W_\sigma &= \Big\{K_5 \left[c^*_\sigma(f_1) c_\sigma(\varepsilon f_1) + \text{h.c.}\right] + K_6 c^*_\sigma(f_1) c_\sigma(f_1)\Big\} \\
&\quad + \Big\{K_2 c^*_\sigma(f_1) c_\sigma(f_1) + K_3 c^*_\sigma(F_2) c_\sigma(F_2) \\
&\qquad + K_4 \left[c^*_\sigma(f_1) c_\sigma(F_2) + \text{h.c.}\right] \Big\} n_d
\end{aligned} \tag{86}$$

$$\begin{aligned}
W_{\text{sd}} &= K_2 \vec{S}_c(f_1 \otimes f_1) . \vec{S}_d + K_3 \vec{S}_c(F_2 \otimes F_2) . \vec{S}_d \\
&\quad + K_4 \Big[\vec{S}_c(f_1 \otimes F_2) + \vec{S}_c(F_2 \otimes f_1)\Big] . \vec{S}_d
\end{aligned} \tag{87}$$

$$F_2 = (\varepsilon + \mu_2)^{-1} f_1 \tag{88}$$

The various coefficients have the following expression

$$\varepsilon_R = -\lambda_1 \tag{89}$$

$$\begin{aligned}
U_R &= \frac{U}{(1+x_1^2)^2} \left[1 - 2\sqrt{2} x_1 \alpha_2 \beta_2 \omega_{12} - \beta_2^2 (1 - 2x_1^2 \omega_{12}^2)\right] \\
&\quad + \beta_2^2 \left\{\lambda_1 - \mu_2 + \omega_{12}^2 \left[x_1^2(\lambda_1 - \mu_1) + (\sqrt{1+x_1^2} - 1)(\mu_1 - \mu_2)\right]\right\}
\end{aligned} \tag{90}$$

$$K_1 = \frac{x_1^2(\alpha_2 - \sqrt{2} x_1 \beta_2 \omega_{12})}{(1+x_1^2)^2} U \tag{91}$$

$$K_2 = \frac{\omega_{12}^2}{4}(1-\alpha_2) KC_2 + \frac{U}{4(1+x_1^2)^{5/2}} KU_2 \tag{92}$$

$$\begin{aligned}
KC_2 &= -x_1^2 \Big\{\lambda_1 \left[1 + \alpha_2 - x_1^2 \omega_{12}^2 (1-\alpha_2)\right] - \mu_1 \left[2 - \omega_{12}^2 (2 + x_1^2)(1-\alpha_2)\right] \\
&\qquad + \mu_2 (1-\alpha_2)(1 - 2\omega_{12}^2)\Big\} \\
&\quad + \left(\sqrt{1+x_1^2} - 1\right) \Big\{2\lambda_1 \left[1 + \alpha_2 + x_1^2 - x_1^2 \omega_{12}^2 (1-\alpha_2)\right] \\
&\qquad - \mu_1 \left[4 + 2x_1^2 - 4\omega_{12}^2 (1-\alpha_2) - 3x_1^2 \omega_{12}^2 (1-\alpha_2)\right] \\
&\qquad + \mu_2 (1-\alpha_2) \left[2 - (4 + x_1^2)\omega_{12}^2\right] \Big\}
\end{aligned} \tag{93}$$

$$\begin{aligned}
KU_2 &= x_1^2 \Big\{2 - \beta_2^2 \omega_{12}^2 - 2\sqrt{2} \beta_2 x_1 \omega_{12} \left[1 - (1-\alpha_2)\omega_{12}^2\right] \\
&\qquad + 2x_1^2 \omega_{12}^2 (1-\alpha_2) \left[2 - (1-\alpha_2)\omega_{12}^2\right] \Big\} \\
&\quad + \left(\sqrt{1+x_1^2} - 1\right) \Big\{2\sqrt{2} \beta_2 x_1 \omega_{12} \left[1 - 2(1-\alpha_2)\omega_{12}^2\right] \\
&\qquad + x_1^2 \left[2 - (4 - 4\alpha_2 - \beta_2^2)\omega_{12}^2 + 4\omega_{12}^4 (1-\alpha_2)^2\right] + 2\beta_2^2 \omega_{12}^2 \\
&\qquad - 2\sqrt{2} \beta_2 x_1^3 \omega_{12}^3 (1-\alpha_2) + 2x_1^4 \omega_{12}^4 (1-\alpha_2)^2 \Big\}
\end{aligned} \tag{94}$$

$$\begin{aligned} K_3 &= \frac{U\omega_{12}^2}{4(1+x_1^2)^2}(1-\alpha_2)(\mu_1-\mu_2)^2(2+x_1^2-2\sqrt{1+x_1^2}) \\ &\quad \left[1+\alpha_2-2\sqrt{2}\beta_2 x_1\omega_{12}+2x_1^2\omega_{12}^2(1-\alpha_2)\right] \\ &\quad -\frac{\omega_{12}^2}{4}(1-\alpha_2)(\mu_1-\mu_2)^2 KC_3 \end{aligned} \tag{95}$$

$$\begin{aligned} KC_3 &= \lambda_1(2+x_1^2-2\sqrt{1+x_1^2})\left[1+\alpha_2-x_1^2\omega_{12}^2(1-\alpha_2)\right] \\ &\quad +\mu_1\omega_{12}^2(1-\alpha_2)\left[4+x_1^4-4\sqrt{1+x_1^2}+x_1^2\left(5-3\sqrt{1+x_1^2}\right)\right] \\ &\quad -\mu_2\Big\{2(\sqrt{1+x_1^2}-1)\left[1+\alpha_2+2\omega_{12}^2(1-\alpha_2)\right] \\ &\qquad +x_1^2\left[1+\alpha_2+\omega_{12}^2(1-\alpha_2)(3-\sqrt{1+x_1^2})\right]\Big\} \end{aligned} \tag{96}$$

$$K_4 = -\frac{\omega_{12}^2}{4}(1-\alpha_2)(\mu_2-\mu_1)KC_4+\frac{U\omega_{12}(\mu_2-\mu_1)}{4(1+x_1^2)^2}KU_4 \tag{97}$$

$$\begin{aligned} KC_4 &= \lambda_1\Big\{2\left(\sqrt{1+x_1^2}-1\right)(1+\alpha_2)+x_1^4\omega_{12}^2(1-\alpha_2) \\ &\qquad -x_1^2\left[2+\alpha_2-\omega_{12}^2\sqrt{1+x_1^2}+2\omega_{12}^2(1-\alpha_2)\left(\sqrt{1+x_1^2}-1\right)\right]\Big\} \\ &\quad +\mu_1\Big\{2(1+x_1^2)-\omega_{12}^2(1-\alpha_2)(4+5x_1^2+x_1^4) \\ &\qquad -\sqrt{1+x_1^2}\left[2+x_1^2-\omega_{12}^2(1-\alpha_2)(4+3x_1^2)\right]\Big\} \\ &\quad +\mu_2\Big\{\alpha_2(2+x_1^2)+\omega_{12}^2(1-\alpha_2)(4+3x_1^2) \\ &\qquad -\sqrt{1+x_1^2}\left[2\alpha_2+\omega_{12}^2(1-\alpha_2)(4+x_1^2)\right]\Big\} \end{aligned} \tag{98}$$

$$\begin{aligned} KU_4 &= -x_1^2\omega_{12}(1-\alpha_2)\left[1+\alpha_2-2\sqrt{2}\beta_2 x_1\omega_{12}+2x_1^2\omega_{12}^2(1-\alpha_2)\right] \\ &\quad +\left(\sqrt{1+x_1^2}-1\right)\Big\{2\beta_2^2\omega_{12}+\sqrt{2}\beta_2 x_1\left[1-4(1-\alpha_2)\omega_{12}^2\right] \\ &\qquad -2x_1^2\omega_{12}(1-\alpha_2)\left[1-2\omega_{12}^2(1-\alpha_2)\right]\Big\} \end{aligned} \tag{99}$$

$$K_5 = \sqrt{1+x_1^2}-1 \tag{100}$$

$$K_6 = \frac{2+x_1^2-2\sqrt{1+x_1^2}}{x_1^2}\varepsilon_d+\frac{2-x_1^2-2(1-x_1^2)\sqrt{1+x_1^2}}{x_1^2}\lambda_1 \tag{101}$$

Then, I have to *diagonalize* the various *quadratic forms*. First I define

$$\begin{aligned}
F_3 &= \varepsilon f_1 - \Gamma_1 f_1 && (102)\\
F_4 &= F_2 - \omega_4 f_1 && (103)\\
\Gamma_1 &= \langle f_1, \varepsilon f_1\rangle \ = \ \frac{\lambda_1 + \varepsilon_d}{x_1^2} - \lambda_1 && (104)\\
\omega_4 &= \langle f_1, F_2\rangle \ = \ \left\langle f_1, (\varepsilon + \mu_2)^{-1} f_1 \right\rangle && (105)\\
&= \frac{1}{x_1^2}(1+x_1^2)(\lambda_1 - \mu_2)\, \frac{U - (1+x_1^2)^2(\lambda_1-\mu_2)}{U + (1+x_1^2)^2(\lambda_1-\mu_2)} && (106)\\
f_i &= \frac{F_i}{\|F_i\|}, \quad i \in \{3,4\} && (107)\\
\|F_3\|^2 &= \frac{1}{x_1^4}\left[g^2 x_1^2 - (\lambda_1 + \varepsilon_d)^2\right] && (108)\\
&= \frac{\langle V, (\varepsilon+\lambda_1)^{-2} V\rangle - \langle V, (\varepsilon+\lambda_1)^{-1} V\rangle^2}{\langle V, (\varepsilon+\lambda_1)^{-2} V\rangle^2} && (109)\\
\|F_4\|^2 &= \|F_2\|^2 - \omega_4^2 && (110)\\
\|F_2\|^2 &= \frac{1 + x_1^2\omega_{12}^2}{\omega_{12}^2(\mu_1-\mu_2)^2\left(\sqrt{1+x_1^2} - 1\right)^2} && (111)
\end{aligned}$$

This allows to rewrite

$$\begin{aligned}
W_\sigma &= \Big\{ a_1 c_\sigma^*(f_1) c_\sigma(f_1) + c_1 \left[c_\sigma^*(f_1) c_\sigma(f_3) + \text{h.c.}\right] \Big\}\\
&\quad + \Big\{ a_2 c_\sigma^*(f_1) c_\sigma(f_1) + b_2 c_\sigma^*(f_4) c_\sigma(F_4)\\
&\qquad + c_2 \left[c_\sigma^*(f_1) c_\sigma(f_4) + \text{h.c.}\right] \Big\} n_d && (112)\\
a_1 &= \lambda_1 + \varepsilon_d && (113)\\
b_1 &= 0 && (114)\\
c_1 &= \frac{1}{x_1^2}\left(\sqrt{1+x_1^2} - 1\right)\sqrt{g^2 x_1^2 - (\lambda_1+\varepsilon_d)^2} && (115)\\
a_2 &= K_2 + K_3\omega_4^2 + 2K_4\omega_4 && (116)\\
b_2 &= K_3 \|F_4\|^2 && (117)\\
c_2 &= \|F_4\|(K_3\omega_4 + K_4) && (118)
\end{aligned}$$

It is then a standard exercise to diagonalize W_σ, and one should note that it allows also to diagonalize the spin-spin interaction. I will note

$$\begin{aligned}
\Delta_i &= (a_i - b_i)^2 + 4c_i^2 && (119)\\
\lambda_\pm &= \frac{1}{2}(a_1 \pm \sqrt{\Delta_1}) && (120)\\
\lambda_{AF} &= \frac{1}{2}\left[\sqrt{\Delta_2} - (a_2 + b_2)\right] && (121)\\
\lambda_F &= -\frac{1}{2}\left[\sqrt{\Delta_2} + (a_2 + b_2)\right] && (122)\\
N_\pm^2 &= 2(\Delta_1 \pm a_1\sqrt{\Delta_1}) && (123)\\
N_{AF}^2 &= 2\left[\Delta_2 - (a_2 - b_2)\sqrt{\Delta_2}\right] && (124)\\
N_F^2 &= 2\left[\Delta_2 + (a_2 - b_2)\sqrt{\Delta_2}\right] && (125)\\
\varphi_\pm &= \frac{1}{N_\pm}\left[(\sqrt{\Delta_1} \pm a_1)f_1 + 2c_1 f_3\right] && (126)\\
\varphi_{AF} &= \frac{1}{N_{AF}}\left\{\left[\sqrt{\Delta_2} - (a_2 - b_2)\right] f_1 - 2c_2 f_4\right\} && (127)\\
\varphi_F &= \frac{1}{N_F}\left\{\left[\sqrt{\Delta_2} + (a_2 - b_2)\right] f_1 + 2c_2 f_4\right\} && (128)
\end{aligned}$$

Then I can finally write H_2 as

$$\begin{aligned}
H_2 = {} & H_0 + \varepsilon_R n_d + U_R n_{d,\uparrow} n_{d,\downarrow} + K_1\left[c_\uparrow^*(f_1)c_\downarrow^*(f_1)d_\downarrow d_\uparrow + \text{h.c.}\right]\\
& + \sum_{\substack{\sigma\in\{\uparrow,\downarrow\}\\ i\in\{+,-\}}} \lambda_i c_\sigma^*(\varphi_i)c_\sigma(\varphi_i) - \sum_{\substack{\sigma\in\{\uparrow,\downarrow\}\\ x\in\{AF,F\}}} \lambda_x c_\sigma^*(\varphi_x)c_\sigma(\varphi_x)n_d\\
& + 4\sum_{x\in\{AF,F\}} \lambda_x \vec{S}_c(\varphi_x \otimes \varphi_x).\vec{S}_d + H_{Irr} && (129)
\end{aligned}$$

5 Asymptotics

I will now evaluate the various coefficients in different regimes, assuming

$$g \ll 1 \tag{130}$$

More precisely, I am interested in

- the *local moment* regime
  - $\varepsilon_d < 0$, $(U + \varepsilon_d) > 0$,
  - $|\varepsilon_d|, (U + \varepsilon_d) \gg g^2 \log g^{-1}$.
- the *intermediate valence* regime
  - $|\varepsilon_d| \ll g^2 \log g^{-1} \ll U$.

I will note respectively $C^{(lm)}$ and $C^{(iv)}$ the value of C in the local moment or intermediate valence regimes. I assume that V is continuous and non-vanishing at the Fermi surface and that ε vanishes linearly (this corresponds to the physical situation) so that the following asymptotics hold

$$\begin{aligned}
\left\langle V, (\varepsilon+\lambda)^{-1} V\right\rangle &\sim a_1 \log \lambda^{-1} && (131)\\
\left\langle V, (\varepsilon+\lambda)^{-2} V\right\rangle &\sim a_2 \lambda^{-1} && (132)\\
\left\langle V, (\varepsilon+\lambda_1)^{-2}(\varepsilon+\mu)^{-1} V\right\rangle &\underset{\mu \ll \lambda_1}{\sim} a_1 \lambda_1^{-2} \log\left(\frac{\lambda_1}{\mu}\right) && (133)\\
\left\langle V, (\varepsilon+\lambda_1)^{-2}(\varepsilon+\mu)^{-2} V\right\rangle &\underset{\mu \ll \lambda_1}{\sim} a_2 \lambda_1^{-2} \mu^{-1} && (134)
\end{aligned}$$

Furthermore, I will assume that $|\varepsilon_d|$ is much smaller than the width of the conduction band (*i.e.* the range of ε) and I choose units so that $|\varepsilon_d| \ll 1$.

- $\lambda_1 = -\varepsilon_R$ is given by equation (33)

$$\begin{aligned}
\lambda_1^{(lm)} &\sim -\varepsilon_d && (135)\\
\lambda_1^{(lm)} + \varepsilon_d &\sim a_1 g^2 \log |\varepsilon_d|^{-1} && (136)\\
\lambda_1^{(iv)} &\sim a_1 g^2 \log g^{-2} && (137)
\end{aligned}$$

- I will need x_1 as expansion parameter

$$\begin{aligned}
x_1^2 &= g^2 \left\langle V, (\varepsilon+\lambda_1)^{-1} V\right\rangle \sim a_2 \frac{g^2}{\lambda_1} && (138)\\
\left[x_1^{(lm)}\right]^2 &\sim a_2 \frac{g^2}{|\varepsilon_d|} \ll 1 && (139)\\
\left[x_1^{(iv)}\right]^2 &\sim \frac{a_2}{a_1 \log g^{-2}} \ll 1 && (140)
\end{aligned}$$

- μ_2 is given by equation (69)

$$\begin{aligned}
\log\left(\frac{\lambda_1}{\mu_2}\right) &\sim a_1^{-1}\left(\frac{U-\lambda_1}{U+\lambda_1}\right)\frac{\lambda_1}{g^2} \gg 1 && (141)\\
\mu_2^{(lm)} &\sim \lambda_1 e^{-a_1^{-1}\left(\frac{U-\lambda_1}{U+\lambda_1}\right)\frac{\lambda_1}{g^2}} = O\left(x_1^\infty, \lambda_1^\infty\right) && (142)\\
\mu_2^{(iv)} &\sim g^2 \lambda_1 \sim a_1 g^4 \log g^{-2} = O\left(x_1^\infty, \lambda_1^2\right) && (143)
\end{aligned}$$

- μ_1 is given by equation (79)

$$\begin{aligned}
\mu_1 &\sim -2\lambda_1\left(\frac{U+\lambda_1}{U-\lambda_1}\right) && (144)\\
\mu_1^{(lm)} &\sim -2|\varepsilon_d|\left(\frac{U+|\varepsilon_d|}{U-|\varepsilon_d|}\right) && (145)\\
\mu_1^{(iv)} &\sim -2\lambda_1 && (146)
\end{aligned}$$

- ω_{12} is also an important expansion parameter, it is given by equation (80)

$$\omega_{12}^2 \sim \frac{1}{a_2}\left(\frac{2\lambda_1}{\mu_1}\right)^2 \frac{\mu_2}{g^2 x_1^2} \tag{147}$$

$$\left[\omega_{12}^{(lm)}\right]^2 \sim \frac{1}{a_2^2}\left(\frac{U-|\varepsilon_d|}{U+|\varepsilon_d|}\right)^2 \frac{|\varepsilon_d|}{g^4}\mu_2^{(lm)} = O\left(x_1^{\infty}, \lambda_1^{\infty}\right) \tag{148}$$

$$\left[\omega_{12}^{(iv)}\right]^2 \sim \frac{a_1^2}{a_2^2} g^2 \left(\log g^{-2}\right)^2 = O(x_1^{\infty}, \lambda_1) \tag{149}$$

- U_R is given by equation (90)

$$U_R = \lambda_1 - \mu_2 + O\left(\frac{\lambda_1^2 x_1^2 \omega_{12}^2}{U}\right) = -\varepsilon_R - \mu_2 + o(\mu_2) \tag{150}$$

- For the scattering potential, I find

$$K_1 \sim -\theta\frac{(2\lambda_1 - \mu_1)}{4\sqrt{2}} x_1^3 \omega_{12} \tag{151}$$

$$\lambda_{\pm} \sim \pm\frac{g x_1}{2} \tag{152}$$

$$\lambda_{\pm}^{(lm)} \sim \pm\frac{\sqrt{a_2}}{2}\frac{g^2}{|\varepsilon_d|^{1/2}} \tag{153}$$

$$\lambda_{\pm}^{(iv)} \sim \pm\frac{1}{2}\sqrt{\frac{a_2}{a_1}}\frac{g}{(\log g^{-2})^{1/2}} \tag{154}$$

$$\varphi_{\pm} \sim \frac{f_1 \pm f_3}{\sqrt{2}} \tag{155}$$

- Finally, for the spin-spin part

$$\lambda_{AF}^{(lm)} \sim \frac{a_2 g^2}{2}\left(\frac{U}{U+\varepsilon_d}\right) > 0 \tag{156}$$

$$\lambda_{F}^{(lm)} \sim -\frac{(U+\varepsilon_d)}{4} < 0 \tag{157}$$

$$\varphi_{AF}^{(lm)} \sim f_1 - \theta\sqrt{2a_2}\frac{g}{|\varepsilon_d|^{1/2}}\left(\frac{U}{U+\varepsilon_d}\right) f_4 \tag{158}$$

$$\varphi_{F}^{(lm)} \sim \sqrt{2a_2}\frac{g}{|\varepsilon_d|^{1/2}}\left(\frac{U}{U+\varepsilon_d}\right) f_1 + \theta f_4 \tag{159}$$

$$\lambda_{AF}^{(iv)} \sim \frac{a_2 g^2}{2} > 0 \tag{160}$$

$$\lambda_{F}^{(iv)} \sim -\frac{U}{4} < 0 \tag{161}$$

$$\varphi_{AF} \sim f_1 - \theta\sqrt{\frac{a_2}{a_1 \log g^{-1}}} f_4 \tag{162}$$

$$\varphi_{F} \sim \sqrt{\frac{a_2}{a_1 \log g^{-1}}} f_1 + \theta f_4 \tag{163}$$

Thus I find a strong ferro-magnetic interaction in one channel and a weak anti-ferromagnetic one in another channel. But one has also to take into account the potential scattering term

$$-\sum_{\substack{\sigma\in\{\uparrow,\downarrow\}\\ x\in\{AF,F\}}} \lambda_x c_\sigma^*(\varphi_x)c_\sigma(\varphi_x)n_d \tag{164}$$

whose effect is to *suppress* the ferro-magnetic channel. Indeed putting an electron in the orbital φ_F with its spin aligned to the impurity one will result in an energy gain (from this part of the interaction)

$$\Delta E_F = -\lambda_F + 4\lambda_F \times \frac{1}{4} = 0 \tag{165}$$

On the other hand, putting an electron in the orbital φ_{AF} with its spin opposite to the impurity one will result in an energy gain

$$\Delta E_{AF} = -\lambda_{AF} + 4\lambda_{AF} \times \left(-\frac{1}{4}\right) = -2\lambda_{AF} < 0 \tag{166}$$

Therefore at low temperature one will see only the anti-ferromagnetic channel and the model will exhibit a Kondo effect.

6 Computation in the general $\mu \neq 0$ case

In the general case, as $\varepsilon - \mu$ is no longer positive one has to go to the so-called *particle-hole* representation. Formally, this amounts to perform the change

$$c_\sigma(k) \longmapsto a_\sigma(k)\theta\left[\varepsilon(k)-\mu\right] + b_\sigma^*(k)\theta\left[\mu-\varepsilon(k)\right] \tag{167}$$

and then do some normal ordering with respect to the a^*'s and b^*'s.

Now one would like to kill the terms which are linear in a_σ, a_σ^*, b_σ or b_σ^*. One can still apply the two general unitary transformations U_1 and U_2 but then one must perform some normal ordering. Thus terms of order 3, 5, *etc.* in c_σ or c_σ^* will give some contribution to the terms linear in the a's or b's.

One can still use my computation of the flow of U_1 and U_2, change the c's into a's and b's, do the normal ordering and set the linear terms to zero. Once again one will have two constraints to meet with two adjustable functions so this should be a priori possible, the only point is that the constraints have now much more complicated expressions.

7 Conclusion

In this paper, I showed that with the help of computer algebra it is possible to compute explicitly non trivial unitary transformations in Quantum Field Theory. For the Anderson impurity model, I succeeded in eliminating the hybridization term in the slightly simplified case of zero chemical potential. This is a rigorous version of the well known Schrieffer-Wolff transformation. Furthermore it should be possible to treat the general case in the same way.

More generally, such exact unitary transformations should be useful when one wants to decouple some small system with finitely many degrees of freedom from a background field, *e.g.* in dissipative systems. I expect also that one could use computer algebra to perform *Hamiltonian conditioning.* For instance, in the case of the Kondo problem, I am investigating the possibility of deriving explicitly the effective low-temperature Hamiltonian for the s-d model. This would enable some Renormalization Group study of the model in the low-temperature phase.

References

[1] A. C. Hewson, *The Kondo Problem to Heavy Fermions*, Cambridge Studies in Magnetism Vol. 2, Cambridge University Press, Cambridge (1993).

[2] S. K. Kehrein and A. Mielke, "Theory of the Anderson impurity model: The Schrieffer-Wolff transformation re-examined", *Ann. Phys.* **252** (1996), 1.

[3] H. R. Krishna-murthy, J. W. Wilkins and K. G. Wilson, *Phys. Rev. B* **21** (1980), 1003 and 1044.

[4] J. R. Schrieffer and P. A. Wolff, *Phys. Rev.* **149** (1966), 491.

[5] A. M. Tsvelick and P. B. Wiegmann, *Adv. Phys.* **32** (1983), 453.

A Differential Flow of $O \mapsto U_2(t)\,O\,U_2^*(t)$

The interested reader will find here the raw material to compute the flow of my second unitary transformation. I put only the differential flow since the integrated one would be too long.

$$
\begin{aligned}
P_1 &= (d_\sigma)^*\, d_\sigma \\
P_2 &= a_\sigma[f_2]^*\, a_\sigma[f_2] \\
P_3 &= a_\sigma[f_2]^*\, d_\sigma \\
P_4 &= a_\sigma[f_2]\, (d_\sigma)^* \\
P_5 &= a_\sigma[f_2]^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} \\
P_6 &= a_\sigma[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
P_7 &= (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} \\
P_8 &= a_\sigma[f_2]^*\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^* \\
P_9 &= a_\sigma[f_2]^*\, a_\sigma[f_2]\, (d_{-\sigma})^*\, d_{-\sigma} \\
P_{10} &= a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_{-\sigma} \\
P_{11} &= a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} \\
P_{12} &= a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* \\
P_{13} &= a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma \\
P_{14} &= a_\sigma[f_2]^*\, a_{-\sigma}[f_2]^*\, d_\sigma\, d_{-\sigma} \\
P_{15} &= a_\sigma[f_2]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^* \\
P_{16} &= a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, d_{-\sigma} \\
P_{17} &= a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]\, (d_{-\sigma})^* \\
P_{18} &= a_\sigma[f_2]^*\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, d_\sigma \\
P_{19} &= a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^* \\
P_{20} &= a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* \\
P_{21} &= a_\sigma[f_2]^*\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} \\
P_{22} &= a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} \\
P_{23} &= a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
P_{24} &= a_\sigma[f_2]^*\, a_\sigma[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} \\
P_{25} &= a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma \\
P_{26} &= a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} \\
P_{27} &= a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_{-\sigma})^*\, d_{-\sigma}
\end{aligned}
$$

$$\begin{aligned}
\partial_t P_1 &= P_5 - P_6 \\
\partial_t P_2 &= -P_5 + P_6 \\
\partial_t P_3 &= -P_7 + P_8 + P_9 + P_{14} \\
\partial_t P_4 &= P_7 - P_9 - P_{10} - P_{15} \\
\partial_t P_5 &= -P_7 + P_8 + P_9 \\
\partial_t P_6 &= P_7 - P_9 - P_{10} \\
\partial_t P_7 &= P_5 - P_6 + P_{11} - P_{12} \\
\partial_t P_8 &= -P_5 + P_{12} - P_{20} + P_{21} \\
\partial_t P_9 &= -P_5 + P_6 - P_{20} + P_{22} \\
\partial_t P_{10} &= P_6 - P_{11} + P_{22} - P_{23} \\
\partial_t P_{11} &= -P_7 + P_{10} + P_{13} \\
\partial_t P_{12} &= P_7 - P_8 - P_{13} \\
\partial_t P_{13} &= -P_{11} + P_{12} + P_{21} - P_{23} \\
\partial_t P_{14} &= -P_{16} - P_{18} + P_{21} + P_{22} \\
\partial_t P_{15} &= P_{17} + P_{19} - P_{20} - P_{23} \\
\partial_t P_{16} &= P_{14} - P_{24} + P_{25} \\
\partial_t P_{17} &= -P_{15} + P_{24} - P_{25} \\
\partial_t P_{18} &= P_{14} - P_{26} + P_{27} \\
\partial_t P_{19} &= -P_{15} + P_{26} - P_{27} \\
\partial_t P_{20} &= P_{24} - P_{25} \\
\partial_t P_{21} &= -P_{26} + P_{27} \\
\partial_t P_{22} &= -P_{24} + P_{25} \\
\partial_t P_{23} &= P_{26} - P_{27} \\
\partial_t P_{24} &= -P_{20} + P_{22} \\
\partial_t P_{25} &= P_{20} - P_{22} \\
\partial_t P_{26} &= P_{21} - P_{23} \\
\partial_t P_{27} &= -P_{21} + P_{23}
\end{aligned}$$

$$
\begin{aligned}
R_1 &= a_\sigma[f_1]^*\, a_\sigma[f_1] \\
R_2 &= a_\sigma[f_1]^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - a_\sigma[f_1]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_3 &= a_\sigma[f_1]^*\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^* + a_\sigma[f_1]^*\, a_\sigma[f_2]\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, (d_{-\sigma})^*\, d_{-\sigma} + a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_{-\sigma} \\
R_4 &= -a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} \\
R_5 &= a_\sigma[f_1]^*\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} - \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* - \\
&\qquad a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_6 &= -a_\sigma[f_1]^*\, a_\sigma[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma - \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma \\
R_7 &= a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^* + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_{-\sigma} \\
R_8 &= a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_9 &= -a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* \\
R_{10} &= -a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_2]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{11} &= a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{12} &= a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma \\
R_{13} &= a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{14} &= a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{15} &= a_\sigma[f_1]^*\, a_\sigma[\epsilon[f_1]] + a_\sigma[\epsilon[f_1]]^*\, a_\sigma[f_1] \\
R_{16} &= a_\sigma[\epsilon[f_1]]^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - a_\sigma[\epsilon[f_1]]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{17} &= a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_1]]\, (d_{-\sigma})^*\, d_{-\sigma} + a_\sigma[\epsilon[f_1]]^*\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^* + \\
&\qquad a_\sigma[\epsilon[f_1]]^*\, a_\sigma[f_2]\, (d_{-\sigma})^*\, d_{-\sigma} + a_\sigma[\epsilon[f_1]]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_{-\sigma} \\
R_{18} &= a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_1]]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} - \\
&\qquad a_\sigma[\epsilon[f_1]]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*
\end{aligned}
$$

$$
\begin{aligned}
R_{19} &= -a_\sigma[f_2]^*\,a_\sigma[\epsilon[f_1]]\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,d_\sigma\,(d_{-\sigma})^* + \\
&\qquad a_\sigma[\epsilon[f_1]]^*\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_2]\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} + \\
&\qquad a_\sigma[\epsilon[f_1]]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,(d_\sigma)^*\,d_\sigma\,d_{-\sigma} - \\
&\qquad a_\sigma[\epsilon[f_1]]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,(d_{-\sigma})^*\,d_{-\sigma} \\
R_{20} &= a_\sigma[f_2]^*\,a_\sigma[\epsilon[f_1]]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_2]\,d_\sigma\,(d_{-\sigma})^* + \\
&\qquad a_\sigma[f_2]^*\,a_\sigma[\epsilon[f_1]]\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,(d_\sigma)^*\,d_{-\sigma} \\
R_{21} &= a_\sigma[f_2]^*\,a_\sigma[\epsilon[f_1]]\,(d_\sigma)^*\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} + \\
&\qquad a_\sigma[\epsilon[f_1]]^*\,a_\sigma[f_2]\,(d_\sigma)^*\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} \\
R_{22} &= a_\sigma[f_2]^*\,a_\sigma[\epsilon[f_1]]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,d_\sigma + \\
&\qquad a_\sigma[\epsilon[f_1]]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,d_\sigma \\
R_{23} &= a_\sigma[f_2]^*\,a_\sigma[\epsilon[f_1]]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_2]\,(d_{-\sigma})^*\,d_{-\sigma} + \\
&\qquad a_\sigma[\epsilon[f_1]]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_2]\,(d_{-\sigma})^*\,d_{-\sigma} \\
R_{24} &= -a_\sigma[f_2]^*\,a_\sigma[\epsilon[f_1]]^*\,a_\sigma[f_2]\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\,a_\sigma[\epsilon[f_1]]\,a_\sigma[f_2]\,(d_\sigma)^*\,(d_{-\sigma})^*\,d_{-\sigma} \\
R_{25} &= a_\sigma[f_2]^*\,a_\sigma[\epsilon[f_1]]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_2]\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} - \\
&\qquad a_\sigma[f_2]^*\,a_\sigma[\epsilon[f_1]]\,a_\sigma[f_2] \\
&\qquad\quad a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,(d_{-\sigma})^*\,d_{-\sigma} \\
R_{26} &= a_\sigma[\epsilon[f_1]]^*\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_2]\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} - \\
&\qquad a_\sigma[\epsilon[f_1]]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,(d_{-\sigma})^*\,d_{-\sigma} \\
R_{27} &= a_\sigma[f_2]^*\,a_\sigma[\epsilon[f_1]]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} + \\
&\qquad a_\sigma[\epsilon[f_1]]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} \\
R_{28} &= a_\sigma[f_1]^*\,a_\sigma[f_1]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1] \\
R_{29} &= -a_\sigma[f_1]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} - \\
&\qquad a_\sigma[f_1]^*\,a_\sigma[f_1]\,a_{-\sigma}[f_1]^*\,(d_\sigma)^*\,d_\sigma\,d_{-\sigma} + \\
&\qquad a_\sigma[f_1]^*\,a_\sigma[f_1]\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,d_\sigma\,(d_{-\sigma})^* + \\
&\qquad a_\sigma[f_1]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,(d_{-\sigma})^*\,d_{-\sigma} \\
R_{30} &= a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} + \\
&\qquad a_\sigma[f_1]^*\,a_\sigma[f_1]\,(d_\sigma)^*\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} \\
R_{31} &= a_\sigma[f_1]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,d_\sigma\,(d_{-\sigma})^* + \\
&\qquad a_\sigma[f_1]^*\,a_\sigma[f_1]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,d_\sigma + \\
&\qquad a_\sigma[f_1]^*\,a_\sigma[f_1]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,d_\sigma + \\
&\qquad a_\sigma[f_1]^*\,a_\sigma[f_1]\,a_\sigma[f_2]\,a_{-\sigma}[f_1]^*\,(d_\sigma)^*\,d_{-\sigma} + \\
&\qquad a_\sigma[f_1]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,(d_{-\sigma})^*\,d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\,a_\sigma[f_1]^*\,a_\sigma[f_1]\,a_{-\sigma}[f_1]\,d_\sigma\,(d_{-\sigma})^* + \\
&\qquad a_\sigma[f_2]^*\,a_\sigma[f_1]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,(d_{-\sigma})^*\,d_{-\sigma} + \\
&\qquad a_\sigma[f_1]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,d_{-\sigma}
\end{aligned}
$$

$$\begin{aligned}
R_{32} \quad = \quad & -a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* + \\
& \quad a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} + \\
& \quad a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} - \\
& \quad a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* - \\
& \quad a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} + \\
& \quad a_\sigma[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
& \quad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - \\
& \quad a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{33} \quad = \quad & a_\sigma[f_1]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
& \quad a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} - \\
& \quad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* - \\
& \quad a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{34} \quad = \quad & a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} + \\
& \quad a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* - \\
& \quad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - \\
& \quad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} \\
R_{35} \quad = \quad & a_\sigma[f_1]^*\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - \\
& \quad a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} + \\
& \quad a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} + \\
& \quad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
& \quad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} - \\
& \quad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* - \\
& \quad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* - \\
& \quad a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{36} \quad = \quad & a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma + \\
& \quad a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma + \\
& \quad a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_{-\sigma} + \\
& \quad a_\sigma[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^* + \\
& \quad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^* + \\
& \quad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, (d_{-\sigma})^*\, d_{-\sigma} + \\
& \quad a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_{-\sigma})^*\, d_{-\sigma} + \\
& \quad a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_{-\sigma} \\
R_{37} \quad = \quad & a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
& \quad a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma}
\end{aligned}$$

$$
\begin{aligned}
R_{38} &= a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma \\
R_{39} &= a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1] \\
&\qquad\quad a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^* + a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_2] \\
&\qquad\quad a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^* + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_{-\sigma} \\
R_{40} &= a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - \\
&\qquad a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1] \\
&\qquad\quad a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma - a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2] \\
&\qquad\quad a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_{-\sigma})^*\, d_{-\sigma} - a_\sigma[f_2]^*\, a_\sigma[f_1]^* \\
&\qquad\quad a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma \\
R_{41} &= a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{42} &= a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^* \\
&\qquad\quad a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^* \\
&\qquad\quad a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{43} &= -a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{44} &= a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} - \\
&\qquad a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* + \\
&\qquad a_\sigma[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - \\
&\qquad a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma}
\end{aligned}
$$

$$\begin{array}{rcl}
R_{45} & = & a_\sigma[f_2]^*\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
 & & \quad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} - \\
 & & \quad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* - \\
 & & \quad a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{46} & = & a_\sigma[f_2]^*\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - \\
 & & \quad a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{47} & = & a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} + \\
 & & \quad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - \\
 & & \quad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* - \\
 & & \quad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{48} & = & -a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} + \\
 & & \quad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - \\
 & & \quad a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* + \\
 & & \quad a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} \\
R_{49} & = & -a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} + \\
 & & \quad a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* - \\
 & & \quad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
 & & \quad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{50} & = & a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^* \\
 & & \qquad a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} - a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1] \\
 & & \qquad a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* + \\
 & & \quad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1] \\
 & & \qquad a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_\sigma[f_2] \\
 & & \qquad a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{51} & = & -a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} + \\
 & & \quad a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} + \\
 & & \quad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - \\
 & & \quad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* \\
R_{52} & = & a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* - \\
 & & \quad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - \\
 & & \quad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} + \\
 & & \quad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{53} & = & a_\sigma[f_2]^*\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^* + \\
 & & \quad a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_{-\sigma})^*\, d_{-\sigma} + \\
 & & \quad a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, (d_{-\sigma})^*\, d_{-\sigma} + \\
 & & \quad a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_{-\sigma}
\end{array}$$

$$
\begin{aligned}
R_{54} &= a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{55} &= a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^* \\
&\qquad\quad a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{56} &= a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{57} &= a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma \\
R_{58} &= a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{59} &= a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{60} &= a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2] \\
&\qquad\quad (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1] \\
&\qquad\quad a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1] \\
&\qquad\quad (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^* \\
&\qquad\quad a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{61} &= a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2] \\
&\qquad\quad (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^* \\
&\qquad\quad a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{62} &= a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^* + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_{-\sigma} \\
R_{63} &= a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} - \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* \\
R_{64} &= -a_\sigma[f_1]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, d_\sigma + a_\sigma[f_1]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^* \\
R_{65} &= a_\sigma[f_1]^*\, a_{-\sigma}[f_1]^*\, d_\sigma\, d_{-\sigma} + a_\sigma[f_1]\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, (d_{-\sigma})^* \\
R_{66} &= a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{67} &= a_\sigma[f_1]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^* + \\
&\qquad a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, (d_\sigma)^*\, d_{-\sigma} \\
R_{68} &= a_\sigma[f_1]^*\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, d_\sigma\, d_{-\sigma} + \\
&\qquad a_\sigma[f_1]\, a_{-\sigma}[f_1]^*\, a_{-\sigma}[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*
\end{aligned}
$$

$$\begin{aligned}
R_{69} &= -a_\sigma[f_1]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_2]\,d_\sigma - \\
&\qquad a_\sigma[f_1]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_1]^*\,d_{-\sigma} + a_\sigma[f_2]^*\,a_\sigma[f_1]\,a_{-\sigma}[f_1]\,(d_{-\sigma})^* + \\
&\qquad a_\sigma[f_1]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]\,(d_\sigma)^*
\end{aligned}$$

$$\begin{aligned}
R_{70} &= -a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,d_\sigma\,(d_{-\sigma})^* + \\
&\qquad a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,d_\sigma\,d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} - \\
&\qquad a_\sigma[f_2]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,(d_{-\sigma})^*\,d_{-\sigma}
\end{aligned}$$

$$\begin{aligned}
R_{71} &= a_\sigma[f_1]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} - \\
&\qquad a_\sigma[f_1]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,(d_{-\sigma})^*\,d_{-\sigma}
\end{aligned}$$

$$\begin{aligned}
R_{72} &= a_\sigma[f_1]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_2]\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} - \\
&\qquad a_\sigma[f_1]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,(d_{-\sigma})^*\,d_{-\sigma}
\end{aligned}$$

$$\begin{aligned}
R_{73} &= -a_\sigma[f_1]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,d_\sigma\,(d_{-\sigma})^* + \\
&\qquad a_\sigma[f_2]^*\,a_\sigma[f_1]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,d_\sigma\,d_{-\sigma}
\end{aligned}$$

$$\begin{aligned}
R_{74} &= a_\sigma[f_1]^*\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} + \\
&\qquad a_\sigma[f_1]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,d_\sigma\,d_{-\sigma} - \\
&\qquad a_\sigma[f_2]^*\,a_\sigma[f_1]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,d_\sigma\,(d_{-\sigma})^* - \\
&\qquad a_\sigma[f_1]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,(d_{-\sigma})^*\,d_{-\sigma}
\end{aligned}$$

$$\begin{aligned}
R_{75} &= -a_\sigma[f_1]^*\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,d_\sigma - \\
&\qquad a_\sigma[f_1]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\,a_\sigma[f_1]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,(d_{-\sigma})^* + \\
&\qquad a_\sigma[f_1]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,(d_\sigma)^*
\end{aligned}$$

$$\begin{aligned}
R_{76} &= a_\sigma[f_1]^*\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_2]\,d_\sigma\,d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\,a_\sigma[f_1]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_1]^*\,d_\sigma\,d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\,a_\sigma[f_1]\,a_\sigma[f_2]\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,(d_{-\sigma})^* + \\
&\qquad a_\sigma[f_1]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,(d_{-\sigma})^*
\end{aligned}$$

$$\begin{aligned}
R_{77} &= a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,d_\sigma + \\
&\qquad a_\sigma[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,d_\sigma\,(d_{-\sigma})^* + \\
&\qquad a_\sigma[f_2]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,(d_{-\sigma})^*\,d_{-\sigma} + \\
&\qquad a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,d_{-\sigma}
\end{aligned}$$

$$\begin{aligned}
R_{78} &= a_\sigma[f_2]^*\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} - \\
&\qquad a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,(d_{-\sigma})^*\,d_{-\sigma}
\end{aligned}$$

$$\begin{aligned}
R_{79} &= -a_\sigma[f_2]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,d_\sigma\,(d_{-\sigma})^* + \\
&\qquad a_\sigma[f_2]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,d_\sigma\,d_{-\sigma}
\end{aligned}$$

$$\begin{aligned}
R_{80} &= a_\sigma[f_1]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_2]\,(d_{-\sigma})^*\,d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\,a_\sigma[f_1]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]\,(d_{-\sigma})^*\,d_{-\sigma}
\end{aligned}$$

$$\begin{aligned}
R_{81} &= a_\sigma[f_2]^*\,a_\sigma[f_1]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,d_\sigma\,(d_{-\sigma})^* + \\
&\qquad a_\sigma[f_2]^*\,a_\sigma[f_1]\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,d_{-\sigma}
\end{aligned}$$

$$
\begin{aligned}
R_{82} &= a_\sigma[f_1]^* \, a_\sigma[f_2] \, a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_1] \, a_{-\sigma}[f_2] \, (d_\sigma)^* \, d_\sigma + \\
&\qquad a_\sigma[f_2]^* \, a_\sigma[f_1] \, a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_1] \, a_{-\sigma}[f_2] \, (d_\sigma)^* \, d_\sigma \\
R_{83} &= a_\sigma[f_1]^* \, a_\sigma[f_2] \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_1] \, (d_\sigma)^* \, d_\sigma \, (d_{-\sigma})^* \, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^* \, a_\sigma[f_1] \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_1] \, (d_\sigma)^* \, d_\sigma \, (d_{-\sigma})^* \, d_{-\sigma} \\
R_{84} &= a_\sigma[f_1]^* \, a_\sigma[f_2] \, a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1]^* \\
&\qquad\quad a_{-\sigma}[f_1] \, a_{-\sigma}[f_2] \, (d_{-\sigma})^* \, d_{-\sigma} + a_\sigma[f_2]^* \, a_\sigma[f_1] \\
&\qquad\quad a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_1] \, a_{-\sigma}[f_2] \, (d_{-\sigma})^* \, d_{-\sigma} \\
R_{85} &= -a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_1] \, a_{-\sigma}[f_2] \, (d_\sigma)^* \, d_\sigma \, (d_{-\sigma})^* \, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^* \, a_\sigma[f_1]^* \, a_\sigma[f_1] \, a_\sigma[f_2] \, (d_\sigma)^* \, d_\sigma \, (d_{-\sigma})^* \, d_{-\sigma} \\
R_{86} &= a_\sigma[f_1]^* \, a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_2] \, d_\sigma \, d_{-\sigma} + \\
&\qquad a_\sigma[f_1] \, a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1] \, a_{-\sigma}[f_2] \, (d_\sigma)^* \, (d_{-\sigma})^* \\
R_{87} &= a_\sigma[f_2]^* \, a_\sigma[f_1]^* \, a_\sigma[f_2] \, a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_1] \, d_\sigma \, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^* \, a_\sigma[f_1] \, a_\sigma[f_2] \, a_{-\sigma}[f_1]^* \\
&\qquad\quad a_{-\sigma}[f_1] \, a_{-\sigma}[f_2] \, (d_\sigma)^* \, (d_{-\sigma})^* \\
R_{88} &= -a_\sigma[f_1]^* \, a_\sigma[f_2] \, a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_2] \, d_{-\sigma} - \\
&\qquad a_\sigma[f_2]^* \, a_\sigma[f_1]^* \, a_\sigma[f_2] \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_2] \, d_\sigma + \\
&\qquad a_\sigma[f_2]^* \, a_\sigma[f_1] \, a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1] \, a_{-\sigma}[f_2] \, (d_{-\sigma})^* + \\
&\qquad a_\sigma[f_2]^* \, a_\sigma[f_1] \, a_\sigma[f_2] \, a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1] \, (d_\sigma)^* \\
R_{89} &= a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_1] \, a_{-\sigma}[f_2] \, (d_\sigma)^* \, d_\sigma \, (d_{-\sigma})^* \, d_{-\sigma} - \\
&\qquad a_\sigma[f_2]^* \, a_\sigma[f_2] \, a_{-\sigma}[f_2]^* \\
&\qquad\quad a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_1] \, a_{-\sigma}[f_2] \, (d_{-\sigma})^* \, d_{-\sigma} \\
R_{90} &= -a_\sigma[f_2]^* \, a_\sigma[f_2] \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_1] \, (d_\sigma)^* \, d_\sigma \, (d_{-\sigma})^* \, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^* \, a_\sigma[f_2] \, a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_1] \, a_{-\sigma}[f_2] \, (d_\sigma)^* \, d_\sigma \\
R_{91} &= -a_\sigma[f_1]^* \, a_\sigma[f_2] \, a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_2] \, (d_\sigma)^* \, d_\sigma \, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^* \, a_\sigma[f_1] \, a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1] \, a_{-\sigma}[f_2] \, (d_\sigma)^* \, d_\sigma \, (d_{-\sigma})^* \\
R_{92} &= a_\sigma[f_2]^* \, a_\sigma[f_1]^* \, a_\sigma[f_2] \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_1] \, d_\sigma \, (d_{-\sigma})^* \, d_{-\sigma} - \\
&\qquad a_\sigma[f_2]^* \, a_\sigma[f_1] \, a_\sigma[f_2] \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_1] \, (d_\sigma)^* \, (d_{-\sigma})^* \, d_{-\sigma} \\
R_{93} &= a_\sigma[f_2]^* \, a_\sigma[f_1]^* \, a_\sigma[f_2] \, a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_1] \\
&\qquad\quad a_{-\sigma}[f_2] \, d_\sigma \, (d_{-\sigma})^* \, d_{-\sigma} - a_\sigma[f_2]^* \, a_\sigma[f_1] \, a_\sigma[f_2] \\
&\qquad\quad a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_1] \, a_{-\sigma}[f_2] \, (d_\sigma)^* \, (d_{-\sigma})^* \, d_{-\sigma} \\
R_{94} &= -a_\sigma[f_1]^* \, a_\sigma[f_2] \, a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_1] \, (d_\sigma)^* \, d_\sigma \, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^* \, a_\sigma[f_1] \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_1] \, a_{-\sigma}[f_2] \, (d_\sigma)^* \, d_\sigma \, (d_{-\sigma})^* \\
R_{95} &= -a_\sigma[f_1]^* \, a_\sigma[f_2] \, a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1] \, a_{-\sigma}[f_2] \, (d_\sigma)^* \, d_\sigma \, (d_{-\sigma})^* + \\
&\qquad a_\sigma[f_2]^* \, a_\sigma[f_1] \, a_{-\sigma}[f_2]^* \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_2] \, (d_\sigma)^* \, d_\sigma \, d_{-\sigma} \\
R_{96} &= a_\sigma[f_2]^* \, a_\sigma[f_2] \, a_{-\sigma}[f_1]^* \, a_{-\sigma}[f_1] \, (d_\sigma)^* \, d_\sigma \, (d_{-\sigma})^* \, d_{-\sigma}
\end{aligned}
$$

$$
\begin{aligned}
R_{97} &= a_\sigma[f_1]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2] \\
&\qquad (d_\sigma)^*\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} + a_\sigma[f_2]^*\,a_\sigma[f_1]\,a_{-\sigma}[f_2]^* \\
&\qquad a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} \\
R_{98} &= a_\sigma[f_2]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^* \\
&\quad a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} \\
R_{99} &= -a_\sigma[f_1]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_2]\,d_{-\sigma} + \\
&\quad a_\sigma[f_2]^*\,a_\sigma[f_1]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,(d_{-\sigma})^* \\
R_{100} &= a_\sigma[f_2]^*\,a_\sigma[f_1]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^* \\
&\qquad a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,d_\sigma - a_\sigma[f_2]^*\,a_\sigma[f_1] \\
&\qquad a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,(d_\sigma)^* \\
R_{101} &= a_\sigma[f_2]^*\,a_\sigma[f_1]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_2]\,d_\sigma\,d_{-\sigma} + \\
&\quad a_\sigma[f_2]^*\,a_\sigma[f_1]\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^* \\
&\qquad a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,(d_{-\sigma})^* \\
R_{102} &= a_\sigma[f_1]^*\,a_{-\sigma}[f_1]\,d_\sigma\,(d_{-\sigma})^* + a_\sigma[f_1]\,a_{-\sigma}[f_1]^*\,(d_\sigma)^*\,d_{-\sigma} \\
R_{103} &= a_{-\sigma}[f_1]^*\,(d_\sigma)^*\,d_\sigma\,d_{-\sigma} - a_{-\sigma}[f_1]\,(d_\sigma)^*\,d_\sigma\,(d_{-\sigma})^* \\
R_{104} &= a_\sigma[f_1]^*\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} - \\
&\quad a_\sigma[f_1]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,d_\sigma\,(d_{-\sigma})^* + \\
&\quad a_\sigma[f_2]^*\,a_\sigma[f_1]\,a_{-\sigma}[f_1]^*\,(d_\sigma)^*\,d_\sigma\,d_{-\sigma} - \\
&\quad a_\sigma[f_1]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,(d_{-\sigma})^*\,d_{-\sigma} \\
R_{105} &= a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,d_\sigma + a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,d_\sigma + \\
&\quad a_\sigma[f_2]^*\,a_{-\sigma}[f_1]\,d_\sigma\,(d_{-\sigma})^* + a_\sigma[f_2]\,a_{-\sigma}[f_1]^*\,(d_\sigma)^*\,d_{-\sigma} \\
R_{106} &= a_\sigma[f_1]^*\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]\,a_{-\sigma}[f_2]\,d_\sigma\,(d_{-\sigma})^* + \\
&\quad a_\sigma[f_1]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]\,(d_{-\sigma})^*\,d_{-\sigma} + \\
&\quad a_\sigma[f_1]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,d_\sigma + \\
&\quad a_\sigma[f_2]^*\,a_\sigma[f_1]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_1]\,d_\sigma\,(d_{-\sigma})^* + \\
&\quad a_\sigma[f_2]^*\,a_\sigma[f_1]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_2]\,(d_{-\sigma})^*\,d_{-\sigma} + \\
&\quad a_\sigma[f_2]^*\,a_\sigma[f_1]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,d_\sigma + \\
&\quad a_\sigma[f_2]^*\,a_\sigma[f_1]\,a_\sigma[f_2]\,a_{-\sigma}[f_1]^*\,(d_\sigma)^*\,d_{-\sigma} + \\
&\quad a_\sigma[f_1]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,d_{-\sigma} \\
R_{107} &= a_\sigma[f_2]^*\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_2]\,d_\sigma\,(d_{-\sigma})^*\,d_{-\sigma} - \\
&\quad a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,(d_{-\sigma})^*\,d_{-\sigma} \\
R_{108} &= a_\sigma[f_2]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_1]^*\,(d_\sigma)^*\,d_\sigma\,d_{-\sigma} - \\
&\quad a_\sigma[f_2]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,d_\sigma\,(d_{-\sigma})^* \\
R_{109} &= a_\sigma[f_2]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_2]\,(d_{-\sigma})^*\,d_{-\sigma} + \\
&\quad a_\sigma[f_2]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]\,(d_{-\sigma})^*\,d_{-\sigma} \\
R_{110} &= a_\sigma[f_2]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_1]^*\,a_{-\sigma}[f_2]\,(d_\sigma)^*\,d_\sigma + \\
&\quad a_\sigma[f_2]^*\,a_\sigma[f_2]\,a_{-\sigma}[f_2]^*\,a_{-\sigma}[f_1]\,(d_\sigma)^*\,d_\sigma
\end{aligned}
$$

$$
\begin{aligned}
R_{111} &= a_\sigma[f_1]^*\, a_\sigma[f_1]\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{112} &= a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} - \\
&\qquad a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* \\
R_{113} &= a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^* \\
R_{114} &= a_\sigma[f_1]^*\, a_\sigma[f_1]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - \\
&\qquad a_\sigma[f_1]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma \\
R_{115} &= \int a_\sigma[\delta_k]^*\, a_\sigma[\delta_k]\epsilon[k]\, dk \\
R_{116} &= -a_\sigma[\epsilon[f_2]]^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + a_\sigma[\epsilon[f_2]]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{117} &= a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_2]]\, (d_{-\sigma})^*\, d_{-\sigma} + a_\sigma[\epsilon[f_2]]^*\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^* + \\
&\qquad a_\sigma[\epsilon[f_2]]^*\, a_\sigma[f_2]\, (d_{-\sigma})^*\, d_{-\sigma} + a_\sigma[\epsilon[f_2]]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_{-\sigma} \\
R_{118} &= a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_2]]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} - \\
&\qquad a_\sigma[\epsilon[f_2]]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* \\
R_{119} &= -a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_2]]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* + \\
&\qquad a_\sigma[\epsilon[f_2]]^*\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[\epsilon[f_2]]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} - \\
&\qquad a_\sigma[\epsilon[f_2]]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{120} &= a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_2]]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^* + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_2]]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_{-\sigma} \\
R_{121} &= -a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_2]]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_2]]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma - \\
&\qquad a_\sigma[\epsilon[f_2]]^*\, a_\sigma[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[\epsilon[f_2]]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma \\
R_{122} &= a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_2]]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[\epsilon[f_2]]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{123} &= -a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_2]]^*\, a_\sigma[f_2]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_2]]\, a_\sigma[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_2]]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_2]]\, a_\sigma[f_2] \\
&\qquad\quad a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{124} &= a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_2]]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^* - \\
&\qquad a_\sigma[\epsilon[f_2]]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, (d_\sigma)^*\, d_\sigma\, d_{-\sigma} \\
R_{125} &= a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_2]]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[\epsilon[f_2]]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{126} &= a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_2]]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + \\
&\qquad a_\sigma[\epsilon[f_2]]^*\, a_\sigma[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma}
\end{aligned}
$$

$$\begin{aligned}
R_{127} &= a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_2]]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - \\
&\qquad a_\sigma[f_2]^*\, a_\sigma[\epsilon[f_2]]\, a_\sigma[f_2] \\
&\qquad\quad a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{128} &= a_\sigma[f_1]^*\, d_\sigma - a_\sigma[f_1]\, (d_\sigma)^* \\
R_{129} &= a_\sigma[f_1]^*\, a_{-\sigma}[f_2]^*\, d_\sigma\, d_{-\sigma} + a_\sigma[f_1]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^* \\
R_{130} &= a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_{-\sigma}[f_2]\, (d_{-\sigma})^* - a_\sigma[f_1]^*\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, d_\sigma \\
&\quad -a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, d_{-\sigma} + a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^* \\
&\quad +a_\sigma[f_1]^*\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} - a_\sigma[f_1]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{131} &= a_\sigma[f_2]^*\, a_\sigma[f_1]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} + a_\sigma[f_1]^*\, a_\sigma[f_2]\, (d_\sigma)^*\, d_\sigma\, (d_{-\sigma})^*\, d_{-\sigma} \\
R_{132} &= -a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, d_\sigma + a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, a_{-\sigma}[f_2]\, (d_\sigma)^* \\
R_{133} &= a_\sigma[f_2]^*\, a_\sigma[f_1]^*\, a_\sigma[f_2]\, a_{-\sigma}[f_2]^*\, d_\sigma\, d_{-\sigma} + a_\sigma[f_2]^*\, a_\sigma[f_1]\, a_\sigma[f_2]\, a_{-\sigma}[f_2]\, (d_\sigma)^*\, (d_{-\sigma})^*
\end{aligned}$$

$$\begin{aligned}
\partial_t R_1 &= -R_2\omega_{12} \\
\partial_t R_2 &= (R_3 - 2P_7\omega_{12}) \\
\partial_t R_3 &= (-2R_2 + 2R_4 + R_5 + (-P_5 + P_6 - P_{11})\omega_{12} + P_{12}\omega_{12}) \\
\partial_t R_4 &= (R_6 + R_7) \\
\partial_t R_5 &= (R_6 + R_8 - 2P_{26}\omega_{12}) \\
\partial_t R_6 &= (-2R_4 + 2R_9 + R_{10} + R_{11}) \\
\partial_t R_7 &= (-R_4 + R_{10} + R_{11} - (P_{20} - P_{22})\omega_{12}) \\
\partial_t R_8 &= (-R_5 - R_9 - (P_{21} - P_{23})\omega_{12}) \\
\partial_t R_9 &= -R_6 \\
\partial_t R_{10} &= (-R_6 - R_7 + R_{12} - 2P_{24}\omega_{12}) \\
\partial_t R_{11} &= (-R_{13} + 2R_{14}\omega_{12}) \\
\partial_t R_{12} &= (-R_4 + R_9 + R_{11}) \\
\partial_t R_{13} &= R_{11} \\
\partial_t R_{14} &= 0 \\
\partial_t R_{15} &= (-R_2\Gamma_{12} - R_{16}\omega_{12}) \\
\partial_t R_{16} &= (R_{17} - 2P_7\Gamma_{12}) \\
\partial_t R_{17} &= (-2R_{16} + 2R_{18} + R_{19} + (-P_5 + P_6 - P_{11} + P_{12})\Gamma_{12}) \\
\partial_t R_{18} &= (R_{20} - R_{21} + R_{22}) \\
\partial_t R_{19} &= (-R_{21} + R_{22} + R_{23} - 2P_{26}\Gamma_{12}) \\
\partial_t R_{20} &= (-R_{18} + R_{24} + R_{25} - (P_{20} - P_{22})\Gamma_{12}) \\
\partial_t R_{21} &= (R_{18} + R_{19} - R_{24} - R_{26}) \\
\partial_t R_{22} &= (-R_{18} - R_{19} + R_{25} + R_{26}) \\
\partial_t R_{23} &= (-R_{26} - (P_{21} - P_{23})\Gamma_{12}) \\
\partial_t R_{24} &= (-R_{20} + R_{21} - 2P_{24}\Gamma_{12}) \\
\partial_t R_{25} &= (-R_{27} + 2R_{14}\Gamma_{12})
\end{aligned}$$

$$
\begin{aligned}
\partial_t R_{26} &= (R_{23} - 2P_{26}\Gamma_{12}) \\
\partial_t R_{27} &= R_{25} \\
\partial_t R_{28} &= R_{29}\omega_{12} \\
\partial_t R_{29} &= (-R_{31} + 2R_{30}\omega_{12}) \\
\partial_t R_{30} &= R_{32} \\
\partial_t R_{31} &= (2R_{29} - 2R_{34} + R_{35} - R_{32}\omega_{12} + R_{33}\omega_{12}) \\
\partial_t R_{32} &= (-4R_{30} + 2R_{36} - R_{6}\omega_{12} + R_{12}\omega_{12} + R_{37}\omega_{12}) \\
\partial_t R_{33} &= (R_{38} + R_{6}\omega_{12} - R_{12}\omega_{12} - R_{37}\omega_{12}) \\
\partial_t R_{34} &= (-R_{39} + R_{40} - 2R_{41}\omega_{12}) \\
\partial_t R_{35} &= (-R_{40} + R_{42} + 2R_{43}\omega_{12}) \\
\partial_t R_{36} &= (-2R_{32} + 2R_{45} + 2R_{48} - R_{9}\omega_{12} - R_{10}\omega_{12} + R_{44}\omega_{12}) \\
\partial_t R_{37} &= (R_{44} + R_{46}) \\
\partial_t R_{38} &= (-R_{33} + R_{47} - R_{4}\omega_{12} - R_{46}\omega_{12}) \\
\partial_t R_{39} &= (R_{34} + R_{49} + R_{50} + R_{47}\omega_{12} + R_{48}\omega_{12}) \\
\partial_t R_{40} &= (-2R_{34} - R_{49} - R_{50} + 2R_{51} - R_{47}\omega_{12} + R_{52}\omega_{12}) \\
\partial_t R_{41} &= R_{47} \\
\partial_t R_{42} &= (-R_{35} + R_{51} - R_{45}\omega_{12}) \\
\partial_t R_{43} &= (-R_{45} - R_{52}) \\
\partial_t R_{44} &= (-2R_{37} + R_{53} + 2P_{26}\omega_{12}) \\
\partial_t R_{45} &= (-2R_{54} + 2R_{55}) \\
\partial_t R_{46} &= (-R_{37} + R_{53}) \\
\partial_t R_{47} &= (-2R_{41} + R_{13}\omega_{12} + R_{56}\omega_{12}) \\
\partial_t R_{48} &= (2R_{57} - 2R_{58} + R_{13}\omega_{12} + R_{56}\omega_{12}) \\
\partial_t R_{49} &= (-R_{39} + R_{59} - 2R_{58}\omega_{12}) \\
\partial_t R_{50} &= (-R_{60} + 2R_{61}\omega_{12}) \\
\partial_t R_{51} &= (-R_{40} + 2R_{43}\omega_{12} + 2R_{54}\omega_{12}) \\
\partial_t R_{52} &= (2R_{43} + 2R_{54} - 2R_{62} - R_{13}\omega_{12} - R_{56}\omega_{12}) \\
\partial_t R_{53} &= (-R_{44} - 2R_{46} + 2R_{63} + (P_{21} - P_{23})\omega_{12}) \\
\partial_t R_{54} &= R_{45} \\
\partial_t R_{55} &= -R_{45} \\
\partial_t R_{56} &= R_{63} \\
\partial_t R_{57} &= (-R_{48} + R_{11}\omega_{12} + R_{63}\omega_{12}) \\
\partial_t R_{58} &= R_{48} \\
\partial_t R_{59} &= (-R_{34} - R_{49} + R_{51}) \\
\partial_t R_{60} &= R_{50} \\
\partial_t R_{61} &= 0 \\
\partial_t R_{62} &= (R_{52} + R_{11}\omega_{12} + R_{63}\omega_{12})
\end{aligned}
$$

$$
\begin{aligned}
\partial_t R_{63} &= (-R_{56} + 2R_{14}\omega_{12}) \\
\partial_t R_{64} &= (-R_{67} - R_{68} - R_{65}\omega_{12} + 2R_{66}\omega_{12}) \\
\partial_t R_{65} &= (R_{33} + R_{69}) \\
\partial_t R_{66} &= R_{70} \\
\partial_t R_{67} &= (-2R_{71} + 2R_{73} + R_{74} - R_{70}\omega_{12} + R_{72}\omega_{12}) \\
\partial_t R_{68} &= (R_{74} + R_{75}) \\
\partial_t R_{69} &= (-R_{38} - 2R_{65} - R_{76} - R_{6}\omega_{12} + R_{12}\omega_{12} + R_{37}\omega_{12}) \\
\partial_t R_{70} &= (-4R_{66} + 2R_{77} + R_{37}\omega_{12}) \\
\partial_t R_{71} &= (R_{67} - 2R_{66}\omega_{12}) \\
\partial_t R_{72} &= (R_{80} - R_{37}\omega_{12}) \\
\partial_t R_{73} &= (R_{81} + R_{82} - R_{83} + R_{41}\omega_{12}) \\
\partial_t R_{74} &= (R_{82} - R_{83} + R_{84} + R_{43}\omega_{12} + R_{85}\omega_{12}) \\
\partial_t R_{75} &= (-2R_{68} - R_{82} + R_{83} - R_{84} - R_{87} - R_{43}\omega_{12} - R_{85}\omega_{12} + R_{86}\omega_{12}) \\
\partial_t R_{76} &= (R_{47} + R_{88}) \\
\partial_t R_{77} &= (-2R_{70} + 2R_{78} + 2R_{79} + R_{44}\omega_{12}) \\
\partial_t R_{78} &= -2R_{89} \\
\partial_t R_{79} &= (2R_{90} + R_{56}\omega_{12}) \\
\partial_t R_{80} &= (-R_{72} - R_{91} - R_{46}\omega_{12}) \\
\partial_t R_{81} &= (-R_{73} - R_{92} + R_{93} + R_{47}\omega_{12} + R_{79}\omega_{12} + R_{91}\omega_{12}) \\
\partial_t R_{82} &= (-R_{73} + R_{93} + R_{94} - R_{91}\omega_{12} + R_{95}\omega_{12}) \\
\partial_t R_{83} &= (R_{73} + R_{92} - R_{94}) \\
\partial_t R_{84} &= (-R_{74} - R_{94} - R_{78}\omega_{12}) \\
\partial_t R_{85} &= (R_{45} - 2R_{78}) \\
\partial_t R_{86} &= (-R_{91} + R_{99}) \\
\partial_t R_{87} &= (R_{93} - R_{100}) \\
\partial_t R_{88} &= (2R_{41} - R_{76} - 2R_{101} - R_{13}\omega_{12} - R_{56}\omega_{12}) \\
\partial_t R_{89} &= 2R_{78} \\
\partial_t R_{90} &= (-2R_{79} + R_{63}\omega_{12}) \\
\partial_t R_{91} &= (R_{41} - R_{13}\omega_{12}) \\
\partial_t R_{92} &= (R_{81} - R_{83} + 2R_{96}\omega_{12}) \\
\partial_t R_{93} &= (-R_{97} + 2R_{98}\omega_{12}) \\
\partial_t R_{94} &= (-R_{82} + R_{83} - R_{43}\omega_{12} - R_{54}\omega_{12}) \\
\partial_t R_{95} &= (-R_{43} - R_{54} + R_{62} + R_{13}\omega_{12}) \\
\partial_t R_{96} &= R_{79} \\
\partial_t R_{97} &= R_{93} \\
\partial_t R_{98} &= 0 \\
\partial_t R_{99} &= (R_{41} - R_{86} - R_{101} - R_{13}\omega_{12})
\end{aligned}
$$

$$
\begin{aligned}
\partial_t R_{100} &= (R_{87} - R_{97} + 2R_{98}\omega_{12} - R_{101}\omega_{12}) \\
\partial_t R_{101} &= 0 \\
\partial_t R_{102} &= (R_{104} - R_{2}\omega_{12} - R_{103}\omega_{12}) \\
\partial_t R_{103} &= (R_{105} - 2P_{7}\omega_{12}) \\
\partial_t R_{104} &= (R_{106} + R_{6}\omega_{12} - R_{12}\omega_{12} - R_{37}\omega_{12}) \\
\partial_t R_{105} &= (2R_{46} - 2R_{103} + R_{107} + R_{108} + (-P_{5} + P_{6} - P_{11} + P_{12})\omega_{12}) \\
\partial_t R_{106} &= (-2R_{52} - 2R_{104} + R_{5}\omega_{12} + 2R_{9}\omega_{12} + R_{10}\omega_{12} - R_{44}\omega_{12} + R_{108}\omega_{12}) \\
\partial_t R_{107} &= (-R_{37} + R_{109}) \\
\partial_t R_{108} &= (R_{110} - 2P_{24}\omega_{12}) \\
\partial_t R_{109} &= (-R_{46} + R_{63} - R_{107}) \\
\partial_t R_{110} &= (-R_{108} + (P_{20} - P_{22})\omega_{12}) \\
\partial_t R_{111} &= (R_{112} - R_{2}\omega_{12}) \\
\partial_t R_{112} &= (R_{113} - 2R_{114}) \\
\partial_t R_{113} &= (-R_{32} + R_{45} + R_{48} + R_{70} - R_{78} - R_{79} - R_{9}\omega_{12}) \\
\partial_t R_{114} &= (R_{32} - R_{48} - R_{70} + R_{79} + R_{112}) \\
\partial_t R_{115} &= R_{116} \\
\partial_t R_{116} &= (-R_{117} + 2P_{7}\Gamma_{2}) \\
\partial_t R_{117} &= (2R_{116} + 2R_{118} + R_{119} + (-P_{5} + P_{6} - P_{11} + P_{12})\Gamma_{2}) \\
\partial_t R_{118} &= (R_{120} + R_{121}) \\
\partial_t R_{119} &= (R_{121} + R_{122} - 2P_{26}\Gamma_{2}) \\
\partial_t R_{120} &= (-R_{118} + R_{123} + (-P_{20} + P_{22})\Gamma_{2}) \\
\partial_t R_{121} &= (-2R_{118} + R_{123} + 2R_{124}) \\
\partial_t R_{122} &= (-R_{119} - R_{124} + (-P_{21} + P_{23})\Gamma_{2}) \\
\partial_t R_{123} &= (-R_{120} - R_{125} + R_{126} - 2(P_{24}\Gamma_{2} - R_{14}\Gamma_{2})) \\
\partial_t R_{124} &= -R_{121} \\
\partial_t R_{125} &= R_{127} \\
\partial_t R_{126} &= (R_{118} - R_{123} - R_{124} + R_{127}) \\
\partial_t R_{127} &= (-R_{125} + 2R_{14}\Gamma_{2}) \\
\partial_t R_{128} &= R_{3} + R_{129} - 2P_{7}\omega_{12} \\
\partial_t R_{129} &= -R_{9} + R_{130} \\
\partial_t R_{130} &= -R_{12} - 2R_{129} + R_{131} - R_{133} \\
\partial_t R_{131} &= R_{4} - R_{9} - R_{10} \\
\partial_t R_{132} &= R_{13} - R_{133} - 2R_{14}\omega_{12} \\
\partial_t R_{133} &= R_{11} + R_{132}
\end{aligned}
$$